



University of Groningen

Expanding diversity in dynamic combinatorial libraries

Leclaire, Julien; Vial, Laurent; Otto, Sijbren; Sanders, Jeremy K.M.

Published in:
Chemical Communications

DOI:
[10.1039/b500638d](https://doi.org/10.1039/b500638d)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2005

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Leclaire, J., Vial, L., Otto, S., & Sanders, J. K. M. (2005). Expanding diversity in dynamic combinatorial libraries: simultaneous exchange of disulfide and thioester linkages. *Chemical Communications*, 1(15). <https://doi.org/10.1039/b500638d>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supporting information for Chemical Communication

Expanding diversity in dynamic combinatorial libraries: simultaneous exchange of disulfide and thioester linkages

Julien Leclaire, Laurent Vial, Sijbren Otto* and Jeremy K. M. Sanders*

NMR ANALYSIS AFTER THE FIRST EQUILIBRIUM

NMR Methods and Parameters

NMR was performed using an Advance 500 Bruker instrument and a standard TCI Cryoprobe ^1H (presaturation).

NMR Spectrum (Figure 4)

1D NOESY with water presaturation during relaxation delay and mixing time (NOESYPR1D).

Solvent: $\text{H}_2\text{O}/\text{CD}_3\text{OD}$ (90/10)

Exact field: 500.13 MHz

Exponential line broadening: 0.30 Hz

Relaxation time (D1): 8.00 seconds

Acquisition time (AQ): 3.172 seconds

Mixing time: 300 milliseconds

Number of points (TD): 65536




Fig. 4

HPLC/ESI-MS ANALYSIS OF THE LIBRARIES

LC-MS Methods and Parameters

LC-MS was performed using an Agilent 1100 series HPLC and Agilent XCT iontrap mass spectrometer. Solvents and formic acid were acquired from Romil.

HPLC Parameters

Injection volume 5µl

Flow rate 1.000 ml/min

Column: Waters symmetry C18 2.1 x 150mm (WAT106005)

Mobile phase: water with 0.1% formic acid (solvent A) and acetonitrile with 0.1% formic acid (solvent B)

Gradient elution

Time (mins)	Solvent B(%)
0	5
26	83
40	95
45	95
50	5
70	5

MS Parameters

Mass range mode: Ultra Scan

Ion polarity: Negative mode

Ion Source: ESI

Dry temperature: 350°C

Nebuliser pressure: 55.00 psi

Dry gas flow: 12 l/min

HV capillary: 4000V

Time (mins)	Target mass:
0 20.3	250
20.3 22.7	450
22.7 24.6	600
24.6 25.5	800
25.5 26.2	1000
26.2 27.9	600
27.9 29.0	1000
29.0 50.9	1100

ICC target: 200000

Base peak chromatogramme (Fig. 5) and extracted ion traces (Fig. 6) of the final library made from 1 after exposure to oxygen.

Structures can be identified and correlated to UV peaks by analysis of the base peak chromatogram. Structures often detected as singly charged species accompanied by the corresponding dimer chelating a sodium ion.




Fig. 5




Fig. 6